

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0

IV PCA

(PATIENT CONTROLLED ANALGESIA)

Fentanyl Hydrochloride 1000_{micrograms}
in 120ml Sodium Chloride 0.9%
(Concentration 8.3 _{micrograms/ml})

Bolus: Protocol E 10 micrograms

Protocol F 20 micrograms

Lockout Period: 5 mins

No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)

Give: Naloxone Dose: up to 0.4mg IV

- Dilute 0.4mg with 3ml Sodium Chloride 0.9% & give in 0.1mg (1ml) increments

Prescriber to sign

Date

Approved by DTC 28/10/2022V1.0