

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1

IV PCA
(PATIENT CONTROLLED ANALGESIA)
**Morphine Sulphate 100mg in 103mL
Sodium Chloride 0.9%**

Bolus: Protocol I 1mg
Protocol J 500 micrograms
Protocol K 1.5 mg
Protocol L 2 mg

Lockout Period: 5 mins
No other opioid to be given whilst PCA is in progress unless directed by anaesthetist
Ensure that all unused stock is recorded in the Ward CD book before disposal

For Respiratory Depression

(resp. rate <8/min &/or sedation Score ≥2)
Give: Naloxone Dose: up to 400microgrammes IV
Dilute 400microgrammes with 3mL Sodium Chloride 0.9% & give 100microgrammes (1mL) increments

Prescriber to sign

Date
Approved by DTAG 26/04/2023 V0.1